

(19) World Intellectual Property
Organization
International Bureau



(43) International Publication Date:
8 July 2004 (08.07.2004)

PCT

(10) International Publication Number
WO 2004/056726 A1

(51) International Patent Classification⁷: **C07B 59/00**

(74) Agents: HAMMETT, Audrey, Grace, Campbell et al.;
Amersham plc, Amersham Place, Little Chalfont, Bucking-
hamshire HP7 9NA (GB).

(21) International Application Number:
PCT/GB2003/005630

(22) International Filing Date:
19 December 2003 (19.12.2003)

(25) Filing Language: English

(26) Publication Language: English

(30) Priority Data:
0229688.7 20 December 2002 (20.12.2002) GB

(81) Designated States (*national*): AE, AG, AL, AM, AT, AU,
AZ, BA, BB, BG, BR, BW, BY, BZ, CA, CH, CN, CO, CR,
CU, CZ, DE, DK, DM, DZ, EC, EE, EG, ES, FI, GB, GD,
GE, GH, GM, HR, HU, ID, IL, IN, IS, JP, KE, KG, KP, KR,
KZ, LC, LK, LR, LS, LT, LU, LV, MA, MD, MG, MK, MN,
MW, MX, MZ, NI, NO, NZ, OM, PG, PH, PL, PT, RO, RU,
SC, SD, SE, SG, SK, SL, SY, TJ, TM, TN, TR, TT, TZ, UA,
UG, US, UZ, VC, VN, YU, ZA, ZM, ZW.

(84) Designated States (*regional*): ARIPO patent (BW, GH,
GM, KE, LS, MW, MZ, SD, SL, SZ, TZ, UG, ZM, ZW),
Eurasian patent (AM, AZ, BY, KG, KZ, MD, RU, TJ, TM),
European patent (AT, BE, BG, CH, CY, CZ, DE, DK, EE,
ES, FI, FR, GB, GR, HU, IE, IT, LU, MC, NL, PT, RO, SE,
SI, SK, TR), OAPI patent (BF, BJ, CF, CG, CI, CM, GA,
GN, GQ, GW, ML, MR, NE, SN, TD, TG).

(71) Applicant (*for all designated States except US*): HAM-
MERSMITH IMANET LIMITED [GB/GB]; Cyclotron
Building, Hammersmith Hospital, Du Cane Road, London
W12 0NN (GB).

(72) Inventors; and

(75) Inventors/Applicants (*for US only*): BRADY, Frank
[GB/GB]; Hammersmith Imanet Limited, Cyclotron
Building, Hammersmith Hospital, Du Cane Road, London
W12 0NN (GB). LUTHRA, Sajinder, Kaur [GB/GB];
Hammersmith Imanet Limited, Cyclotron Building, Ham-
mersmith Hospital, Du Cane Road, London W12 0NN
(GB). ZHAO, Yongjun [CN/GB]; Hammersmith Imanet
Limited, Cyclotron Building, Hammersmith Hospital, Du
Cane Road, London W12 0NN (GB).

Published:

- with international search report
- before the expiration of the time limit for amending the
claims and to be republished in the event of receipt of
amendments

*For two-letter codes and other abbreviations, refer to the "Guid-
ance Notes on Codes and Abbreviations" appearing at the begin-
ning of each regular issue of the PCT Gazette.*

(54) Title: SOLID-PHASE PREPARATION OF [¹⁸F]FLUOROHALOALKANES

(57) Abstract: The invention relates to a process for the production of an [¹⁸F]fluorohaloalkane which comprises treatment of a solid support-bound precursor of formula (I): SOLID SUPPORT-LINKER-SO₂-O-(CH₂)_nX (I) wherein n is an integer of from (1) to (7) and X is chloro, bromo or iodo; with ¹⁸F⁻ to produce the [¹⁸F]fluorohaloalkane of formula (II) ¹⁸F-(CH₂)_n-X (II) wherein n and X are as defined for the compound of formula (I), optionally followed by (i) removal of excess ¹⁸F⁻, for example by ion-exchange chromatography; and/or(ii) removal of organic solvent.



THIS PAGE BLANK (USP 10)

SOLID-PHASE PREPARATION OF [^{18}F]FLUOROHALOALKANES

The present invention relates to a process for the preparation of fluorohaloalkane compounds such as [^{18}F]bromofluoromethane. [^{18}F]Fluorohaloalkanes are important reagents for performing O-, N-, and S- ^{18}F fluoroalkylations and are commonly used to radiolabel radioligands for use in positron emission tomography (PET) studies.

[^{18}F]Fluorohaloalkanes have previously been prepared by nucleophilic displacement, by [^{18}F]F $^-$, of a leaving group from a suitable precursor compound. Thus, for example Zhang *et al*, Applied Radiation and Isotopes 57, 335-342 (2002), describes synthesis of [^{18}F]fluoroethyl bromide by nucleophilic displacement of 2-trifluoromethanesulphonyloxy ethylbromide with [^{18}F]F $^-$ and Seung-Jun *et al*/Applied Radiation and Isotopes (1999), 51, 293-7 describes an analogous synthesis of 3- ^{18}F fluoropropylbromide. A similar method is described in Comagic *et al*/Applied Radiation and Isotopes (2002), 56, 847-851 wherein 2-bromo-1- ^{18}F fluoroethane is prepared by nucleophilic displacement of 1,2-dibromoethane with [^{18}F]F $^-$. Solid-phase nucleophilic fluorination methods are described in co-pending International Patent Application PCT/GB02/02505.


In view of the importance of [^{18}F]fluorohaloalkanes as radiolabelling reagents, there exists the need for synthetic methods for their preparation in good radiochemical yield and in which isolation of the product is more readily achievable. Furthermore, there is also a need for such synthetic methods which are amenable to automation.

In a first aspect, the invention provides a process for the production of an [^{18}F]fluorohaloalkane which comprises treatment of a solid support-bound precursor of formula (I):



wherein n is an integer of from 1 to 7 and X is chloro, bromo, or iodo;

with $^{18}\text{F}^-$ to produce the $[^{18}\text{F}]$ fluorohaloalkane of formula (II)



wherein n and X are as defined for the compound of formula (I), optionally followed by

- (i) removal of excess $^{18}\text{F}^-$, for example by ion-exchange chromatography; and/or
- (ii) removal of organic solvent.


Preferably, in the compounds of formula (I) above, n is an integer of 1 to 4, more preferably, 1 or 2. In the compounds of formula (I) above, X is preferably bromo or iodo. Preferred $[^{18}\text{F}]$ fluorohaloalkanes of formula (II) prepared, include $[^{18}\text{F}]$ fluorobromomethane, $[^{18}\text{F}]$ fluoriodomethane, $[^{18}\text{F}]$ fluorobromoethane, $[^{18}\text{F}]$ fluoriodoethane, $[^{18}\text{F}]$ fluorobromopropane, and $[^{18}\text{F}]$ fluoriodopropane.

As the $[^{18}\text{F}]$ fluorohaloalkane of formula (II) is removed from the solid-phase into solution, all unreacted precursor remains bound to the resin and can be separated by simple filtration, thus obviating the need for complicated purification, for example by HPLC. The $[^{18}\text{F}]$ fluorohaloalkane of formula (II) may be cleaned up by removal of excess F^- , for example by ion-exchange chromatography and/or by removal of any organic solvent.

As shown in Scheme 1, the compound of formula (I) may be conveniently prepared from any sulphonic acid functionalised commercially available resin, such as Merrifield Resin, NovaSyn[®] TG Bromo Resin, (Bromomethyl)phenoxymethyl polystyrene, or Wang Resin which may be reacted with a chlorinating agent to give the corresponding sulphonyl chloride resin. This may be carried out by treating the resin with, for example, phosphorus pentachloride, phosphorus trichloride, oxalyl chloride, or thionyl chloride, in an appropriate inert solvent such as dichloromethane, chloroform, or acetonitrile, and heating at elevated temperature for a period of time. The excess reagent may then be removed from the resin by washing with further portions of the inert solvent. The sulphonyl chloride resin may then be reacted with the alcohol

analogue of the tracer to produce the resin-bound precursor of formula (I). This may be carried out by treating the resin with a solution of the alcohol in an inert solvent such as chloroform, dichloromethane, acetonitrile, or tetrahydrofuran containing a non-nucleophilic soluble base such as sodium hydride or a trialkylamine, for example triethylamine or diisopropylethylamine. The reaction may be carried out at a temperature of 10 to 80°C, optimally at ambient temperature for a period of from around 1 to 24 hours. The excess alcohol and base may then be removed from the solid support by washing with further portions of an inert solvent such as chloroform, dichloromethane, or tetrahydrofuran. Alternatively, the LINKER may be attached to the haloalkane, before being attached to the SOLID SUPPORT to form the compound of formula (I), using analogous chemistry to that described above.


Scheme 1




In the compounds of formula (I), the "SOLID SUPPORT" may be any suitable solid-phase support which is insoluble in any solvents to be used in the process but to which the LINKER and/or haloalkane can be covalently bound. Examples of suitable SOLID SUPPORT include polymers such as polystyrene (which may be block grafted, for example with polyethylene glycol), polyacrylamide, or polypropylene or glass or silicon coated with such a polymer. The solid support may be in the form of small discrete particles such as beads or pins, or as a

coating on the inner surface of a cartridge or on a microfabricated vessel.

In the compounds of formula (I), the "LINKER" may be any suitable organic group which serves to space the reactive site sufficiently from the solid support structure so as to maximise reactivity. Suitably, the LINKER comprises zero to four aryl groups (suitably phenyl) and/or a C₁₋₁₆alkyl (suitably C₁₋₆alkyl) or C₁₋₁₆haloalkyl (suitably C₁₋₆haloalkyl), typically C₁₋₁₆ fluoroalkyl (suitably C₁₋₆ fluoroalkyl), or C₁₋₁₆alkoxy or C₁₋₁₆haloalkoxy (suitably C₁₋₆alkoxy or C₁₋₆haloalkoxy) typically C₁₋₁₆fluoroalkoxy (suitably C₁₋₆fluoroalkoxy), and optionally one to four additional functional groups such as amide or sulphonamide groups. Examples of such linkers are well known to those skilled in the art of solid-phase chemistry, but include:





wherein at each occurrence, k is an integer of 0 to 3, n is an integer of 1 to 16, and R^L is hydrogen or C_{1-6} alkyl.

- 5 Treatment of the compound of formula (I) with $^{18}F^-$ may be effected by treatment with any suitable source of $^{18}F^-$, such as $Na^{18}F$, $K^{18}F$, $Cs^{18}F$, tetraalkylammonium ^{18}F fluoride, or tetraalkylphosphonium ^{18}F fluoride. To increase the reactivity of the fluoride, a phase transfer catalyst such as 4,7,13,16,21,24 hexaoxa-1,10-diazabicyclo[8,8,8] hexacosane may be added and the reaction performed in a
- 10 non protic solvent. These conditions give reactive fluoride ions. The treatment with $^{18}F^-$ is suitably effected in the presence of a suitable organic solvent such as acetonitrile, dimethylformamide, dimethylsulphoxide, tetrahydrofuran, dioxan, 1,2 dimethoxyethane, sulpholane, N-methylpyrrolidine, at a non-extreme temperature, for example, $15^\circ C$ to $180^\circ C$, preferably at elevated temperature.
- 15 On completion of the reaction, the $[^{18}F]$ fluorohaloalkane of formula (II) dissolved in the solvent is conveniently separated from the solid-phase by filtration.

Any excess $^{18}F^-$ may be removed from the solution of $[^{18}F]$ fluorohaloalkane by any suitable means, for example by ion-exchange chromatography or solid phase

- 20 absorbents. Suitable ion-exchange resins include BIO-RAD AG 1-X8 or Waters QMA and suitable solid phase absorbents include alumina. The excess $^{18}F^-$ may be removed using such solid phases at room temperature in aprotic solvents.

Any organic solvent may be removed by any standard method such as by

- 25 evaporation at elevated temperature *in vacuo* or by passing a stream of inert gas such as nitrogen or argon over the solution.

According to a further aspect, the invention provides a process for the production of an $[^{18}F]$ fluorohaloalkane which comprises treatment of a solid support-bound precursor of formula (III):


- 30



wherein n and X are as defined for the compound of formula (I), and Y⁻ is an anion (preferably trifluoromethylsulphonate (triflate) anion or tetraphenyl borate anion);

5

with ¹⁸F⁻ to produce the [¹⁸F]fluorohaloalkane of formula (II)




10 wherein n and X are as defined for the compound of formula (III), optionally followed by

- (i) removal of excess ¹⁸F⁻, for example by ion-exchange chromatography; and/or
- (ii) removal of organic solvent.

15 The compound of formula (III) may be conveniently prepared from a functionalised commercially available resin such as a Merrifield Resin or Wang Resin. Suitably, a hydroxyiodoaryl (such as an iodophenol) containing LINKER group is treated with an inorganic base, such as cesium carbonate and then added to the resin, pre-swollen with an inert solvent, such as N,N-dimethylformamide and allowed to
20 react at elevated temperature, for example 30 to 80°C. Excess reagents may be removed by washing the resin with further inert solvent. The resultant iodophenol functionalised resin may then be treated with a source of acetate anions (such as acetic acid, acetic anhydride, or acetyl chloride) in the presence of an oxidising agent, such as hydrogen peroxide to provide the corresponding diacetoxo-
25 iodophenyl functionalised resin. The diacetoxo-iodophenyl functionalised resin may then be stirred in an inert solvent, such as dichloromethane, in the presence of acid such as hydrochloric acid, trifluoromethane sulphonic acid, or acetic acid at a low temperature, suitably -40°C to 10°C before addition of the fluorohaloalkane, suitably functionalised as a boronic acid or triorgano tin (suitably trialkyl tin)
30 derivative which may be coupled to the resin at a non-extreme temperature. As in previous steps, the desired compound of formula (III) may be separated by filtration and washing with an inert solvent.

In the compound of formula (III), the LINKER is as defined above but suitably comprises an aryl group (suitably phenyl) adjacent to the I^+ . Preferred examples include:



5

wherein each phenyl is optionally substituted by 1 to 4 groups selected from C_{1-6} alkyl and C_{1-6} alkoxy, but is suitably unsubstituted.

- 10 Treatment of a compound of formula (III) with F^- , preferred haloalkanes in formula (III), removal of excess F^- and any organic solvent are all suitably as described for the compounds of formula (I) above.

15 The compounds of formula (I) and (III) are novel and thus form a further aspect of the present invention.

As described above, the advantages of such solid-phase processes for preparation of $[^{18}F]$ fluorohaloalkanes include the relative speed of the process, simplified purification methods and ease of automation- all of which mean that the processes are suitable for preparation of $[^{18}F]$ fluorohaloalkanes which can then be used to prepared ^{18}F -labelled tracers for use in PET. Accordingly, the present invention provides a process for the preparation of a $[^{18}F]$ fluorohaloalkane of

20

formula (II) for use in PET chemistry.

Conveniently, the solid support bound precursor of formula (I) or (III) could be provided as part of a kit to a radiopharmacy, PET centre, or nuclear medicine department. The kit may contain a cartridge which can be plugged into a suitably adapted automated synthesiser. The cartridge may contain, apart from the solid support-bound precursor, a column to remove unwanted fluoride ion, and an appropriate vessel connected so as to allow the reaction mixture to be evaporated and allow the product to be formulated as required. The reagents and solvents and other consumables required for the synthesis may also be included together with a compact disc carrying the software which allows the synthesiser to be operated in a way so as to meet the customers requirements for radioactive concentration, volumes, time of delivery etc.

Conveniently, all components of the kit are disposable to minimise the possibilities of contamination between runs and may be sterile and quality assured.

The invention further provides a radiosynthesis kit for the preparation of an [^{18}F]fluorohaloalkane for use in PET chemistry, which comprises:

- (i) a vessel containing a compound of formula (I) or (III); and
- (ii) means for eluting the vessel with a source of $^{18}\text{F}^-$; and
- (iii) an ion-exchange cartridge for removal of excess $^{18}\text{F}^-$.

The invention further provides a cartridge for a radiosynthesis kit for the preparation of an [^{18}F]fluorohaloalkane for use in PET chemistry which comprises:


- (i) a vessel containing a compound of formula (I) or (III); and
- (ii) means for eluting the vessel with a source of $^{18}\text{F}^-$.

The invention will now be illustrated by way of the following Examples.

Example 1. Synthesis of [^{18}F]-fluorobromomethane

Example 1(i) Preparation of perfluorobutane-1,4-bis-sulphonylchloride

(Following the method of Weiming Qiu and Donald J. Burton
Journal of fluorine chemistry, 60 (1993) 93-100.)




The mixture of 1,4 diiodoperfluorobutane ($\text{I}(\text{CF}_2)_4\text{I}$) (24.14g, 53.2mmol), sodium dithionite $\text{Na}_2\text{S}_2\text{O}_4$ (24g, 117.2mmol) and sodium hydrogen sulphate NaHCO_3 (12.8g, 152.4mmol) in water H_2O (36ml) / Acetonitrile CH_3CN (36ml) was stirred at room temperature for 2 hours. It was filtered, and the filtrate was concentrated under reduced pressure to remove the acetonitrile. To the residue was added H_2O (100ml). The so obtained solution was vigorously stirred and treated with chlorine gas Cl_2 at 0°C until the colour of I_2 disappeared. Dichloromethane CH_2Cl_2 (100ml) was added and the mixture vigorously shaken. The organic phase was separated, and the aqueous phase was extracted with CH_2Cl_2 . The combined organic phase was washed with water H_2O , brine, and dried with sodium sulphate Na_2SO_4 and concentrated to afford a waxy yellow crystalline solid. (15.4g, 74%). Recrystallization from hexane afforded off-white needles of perfluorobutane-1,4-bis-sulphonylchloride.

10

15

20 ^{19}F NMR (CDCl_3 , CFCl_3 reference) δ : -104.4, -119.1.

Example 1(ii) Preparation of perfluorobutane-1,4-bis-sulphonate dipotassium salt




25 To the solution of potassium hydroxide KOH (9.8g, 5eq) in water H_2O (19ml) was added gradually perfluorobutane-1,4-bis-sulphonylchloride (14g, 35mmol) at 85°C - 90°C with stirring. After the addition, the reaction was continued for more 4

hours at the same temperature, and then it was cooled overnight. It was filtered and the solids was washed with a little of cooled water and dried in vacuum to give perfluorobutane-1,4-bis-sulphonate dipotassium salt

5 ^{19}F NMR (CD_3OD , CFCl_3 reference) δ : -114.00, -120.11.

Example 1(iii) Preparation of perfluorobutane-1,4-bis-sulphonic acid

(Following the method described in US patent 4329,478, Fred E. Behr.)



Perfluorobutane-1,4-bis-sulphonate dipotassium salt (15g, 34.2mmol) was dissolved in hot water (100ml). It was added to an ion exchange column of
 15 Amberlyst 15 resin, (40x4cm) which had been previously washed with excess 6N HCl and rinsed with distilled water. The column was then washed slowly with distilled water, and the first 300ml of aqueous solution collected. The solution was concentrated in vacuum and the residue was dried under reduced pressure at 80°C to afford perfluorobutane-1,4-bis-sulphonic acid. (11.0g, 30mmol, 88%)


20 ^1H NMR (CDCl_3) δ : 8.00

^{18}F NMR (CDCl_3 , CFCl_3 reference) δ : -114.7, -121.3.

Example 1(iv) Preparation of perfluorobutane-1,4-bis-sulphonic acid anhydride

(Following the method described in US patent 4329,478, Fred E. Behr.)

25




Perfluorobutane-1,4-bis-sulphonic acid (11.0g, ~30mmol) was mixed with P_2O_5 (40g, ~10eq) and sand. The mixture was heated to 140-180°C and distilled under reduced pressure with dry-ice cooling collector to afford crude product (5.12g). Redistillation gives pure perfluorobutane-1,4-bis-sulphonic acid anhydride.

5

^{18}F NMR ($CDCl_3$, $CFCl_3$ reference) δ : -105.7, -121.8.

Example 1(v) Synthesis of PS – 4-(Benzyl-ethyl-sulfonamide)octafluoro-butane-1-sulfonic acid



10

To a portion of the polystyrene resin (Novabiochem, Novasyn resin) (202mg), which had previously been swollen in dichloromethane (2ml) and then suspended in a further aliquot of dichloromethane (2ml) the perfluorobutyl-1,4-cyclic-sulfonic anhydride (116mg, 5Eq) was added. Following this di-isopropylethyl amine (0.174 ml) was added and the suspension stirred overnight at room temperature. The solvent was removed by filtration and the resin washed with consecutive addition and filtration of dichloromethane (5 ml), methanol (5ml), DMF (5ml), water (5 ml), methanol (5 ml), and dichloromethane (5ml). The resulting resin was then treated with NaOH (1M) in THF/water (2 x 2ml) before washing with consecutive portions of methanol (5ml), dichloromethane (5ml) and methanol (5 ml) again. The resin was then dried under high vacuum.


15

20

Gel Phase ^{19}F NMR (referenced to $CFCl_3$, 300K) : δ -121.0, -114.8, -113.4

Example 1(vi) Synthesis of PS – 4-(Benzyl-ethyl-sulfonamide)octafluoro-butane-1-sulfonyl chloride


25



A portion of the resin prepared in the manner of Example 1(v) above is swollen with dichloromethane (2ml) and then washed consecutively with HCl (1M) in THF/water (10 x 5 ml) to give the free sulphonic acid. The resin is washed
 5 consecutively with dichloromethane, methanol and THF before drying under high vacuum.

The resin is then suspended in dichloromethane and to it is added in excess a common chlorinating agent such as phosphorous pentachloride, phosphorus
 10 trichloride or thionyl chloride. The suspension is stirred for 2 hours before filtration and then washing of the resin with dichloromethane and then THF.

Example 1(vii) Synthesis of fluorobromomethane precursor resin



15 A solution of bromomethanol in THF is added to a portion of the resin prepared as described in Example 1(vi) above which has previously been swollen in THF. To this is added a solution of potassium t-butoxide in tetrahydrofuran and the suspension is stirred overnight. After filtration the resin is washed consecutively with dichloromethane and THF before drying under high vacuum.

20

Example 1(viii) Radiofluorination to prepare [¹⁸F]-fluorobromomethane

To a portion of the resin (prepared as described in Example 1(vii)) held in a

cartridge is added a solution in dry acetonitrile of kryptofix, potassium carbonate and [^{18}F]-fluoride. The suspension is heated to 85°C for 10 minutes and then the solution is filtered off. The solution is then passed onto a C₁₈ solid phase extraction cartridge and washed with water to remove acetonitrile, kryptofix and potassium carbonate. Addition of more acetonitrile washes the radiofluorinated product of the cartridge into a solution of 0.1 M HCl. This solution is heated for 5 minutes before neutralization and analysis.

Claims

1. A process for the production of an [^{18}F]fluorohaloalkane which comprises
5 treatment of a solid support-bound precursor of formula (I):



wherein n is an integer of from 1 to 7 and X is chloro, bromo or iodo;
with $^{18}\text{F}^-$ to produce the [^{18}F]fluorohaloalkane of formula (II)



wherein n and X are as defined for the compound of formula (I), optionally followed by

- (i) removal of excess $^{18}\text{F}^-$, for example by ion-exchange chromatography; and/or
15 (ii) removal of organic solvent.

2. A process for the production of an [^{18}F]fluorohaloalkane according to claim 1 wherein n is an integer of 1 to 4, preferably 1 or 2.

- 20 3. A process for the production of an [^{18}F]fluorohaloalkane according to claim 1 or 2 wherein the compound of formula (II) prepared is selected from [^{18}F]fluorobromomethane, [^{18}F]fluoroiodomethane, [^{18}F]fluorobromoethane, [^{18}F]fluoroiodoethane, [^{18}F]fluorobromopropane, and [^{18}F]fluoroiodopropane

- 25 4. A compound of formula (I) as defined in claim 1:



wherein n is an integer of from 1 to 7 and X is chloro, bromo or iodo.

30

5. A compound of formula (I) according to claim 4 wherein n is an integer of from 1 to 4, and is preferably 1 or 2.

6. A radiosynthesis kit for the preparation of an [^{18}F]fluorohaloalkane for use in PET chemistry, which comprises:

- 5 (i) a vessel containing a compound of formula (I) as defined in any one of claims 1 to 3; and
- (ii) means for eluting the vessel with a source of $^{18}\text{F}^-$; and
- (iii) an ion-exchange cartridge for removal of excess $^{18}\text{F}^-$.

7. A cartridge for a radiosynthesis kit which comprises:

- 10 (i) a vessel containing a compound of formula (I) as defined in any one of claims 1 to 3; and
- (ii) means for eluting the vessel with a source of $^{18}\text{F}^-$.

THIS PAGE BLANK (USPTO)

INTERNATIONAL SEARCH REPORT

International Application No

PCT/GB/05630

A. CLASSIFICATION OF SUBJECT MATTER

IPC 7 C07B59/00

According to International Patent Classification (IPC) or to both national classification and IPC

B. FIELDS SEARCHED

Minimum documentation searched (classification system followed by classification symbols)

IPC 7 C07B

Documentation searched other than minimum documentation to the extent that such documents are included in the fields searched

Electronic data base consulted during the international search (name of data base and, where practical, search terms used)

EPO-Internal, CHEM ABS Data, WPI Data, PAJ, BEILSTEIN Data

C. DOCUMENTS CONSIDERED TO BE RELEVANT

Category *	Citation of document, with indication, where appropriate, of the relevant passages	Relevant to claim No.
P, A	WO 2003/002157 A (AMERSHAM PLC, UK; IMAGING RESEARCH SOLUTIONS LTD.) 9 January 2003 (2003-01-09) claims 1-15; examples 1-3	1-7
A	S. COMAGIC ET. AL.: "EFFICIENT SYNTHESIS OF 2-BROMO-1- ¹⁸ F-FLUOROETHANE AND ITS APPLICATION IN THE AUTOMATED PREPARATION OF ¹⁸ F-FLUORETHYLATED COMPOUNDS" APPLIED RADIATION AND ISOTOPES, vol. 56, 2002, pages 847-851, XP002277733 cited in the application the whole document	1-7

☐ Further documents are listed in the continuation of box C.

☒ Patent family members are listed in annex.

* Special categories of cited documents:

"A" document defining the general state of the art which is not considered to be of particular relevance

"E" earlier document but published on or after the international filing date

"L" document which may throw doubts on priority claim(s) or which is cited to establish the publication date of another citation or other special reason (as specified)

"O" document referring to an oral disclosure, use, exhibition or other means

"P" document published prior to the international filing date but later than the priority date claimed

"T" later document published after the international filing date or priority date and not in conflict with the application but cited to understand the principle or theory underlying the invention

"X" document of particular relevance; the claimed invention cannot be considered novel or cannot be considered to involve an inventive step when the document is taken alone

"Y" document of particular relevance; the claimed invention cannot be considered to involve an inventive step when the document is combined with one or more other such documents, such combination being obvious to a person skilled in the art.

"&" document member of the same patent family

Date of the actual completion of the international search

22 April 2004

Date of mailing of the international search report

04/05/2004

Name and mailing address of the ISA

European Patent Office, P.B. 5818 Patentlaan 2
NL - 2280 HV Rijswijk
Tel. (+31-70) 340-2040, Tx. 31 651 epo nl,
Fax: (+31-70) 340-3016

Authorized officer

Kleidernigg, O

INTERNATIONAL SEARCH REPORT

Information on patent family members

International Application No

PCT/GB 03/05630

Patent document cited in search report	Publication date	Patent family member(s)	Publication date
WO 2003002157 A	09-01-2003	CA 2450637 A1	09-01-2003
		EE 200400008 A	16-02-2004
		WO 03002157 A1	09-01-2003
<hr/>			

**This Page is Inserted by IFW Indexing and Scanning
Operations and is not part of the Official Record**

BEST AVAILABLE IMAGES

Defective images within this document are accurate representations of the original documents submitted by the applicant.

Defects in the images include but are not limited to the items checked:

- ☐ BLACK BORDERS
- ☒ IMAGE CUT OFF AT TOP, BOTTOM OR SIDES
- ☒ FADED TEXT OR DRAWING
- ☐ BLURRED OR ILLEGIBLE TEXT OR DRAWING
- ☐ SKEWED/SLANTED IMAGES
- ☐ COLOR OR BLACK AND WHITE PHOTOGRAPHS
- ☐ GRAY SCALE DOCUMENTS
- ☐ LINES OR MARKS ON ORIGINAL DOCUMENT
- ☐ REFERENCE(S) OR EXHIBIT(S) SUBMITTED ARE POOR QUALITY
- ☐ OTHER: _____

IMAGES ARE BEST AVAILABLE COPY.

As rescanning these documents will not correct the image problems checked, please do not report these problems to the IFW Image Problem Mailbox.

THIS PAGE BLANK (USP 10)